

STRONG EVOLUTION IN THE LUMINOSITY-VELOCITY RELATION AT $Z \gtrsim 1$?

JAMES S. BULLOCK¹

Department of Astronomy, Ohio State University, Columbus, OH 43210

AVISHAI DEKEL²

Racah Institute of Physics, The Hebrew University, Jerusalem 91904, Israel

JOEL R. PRIMACK³

Physics Department, University of California, Santa Cruz, CA 95064

RACHEL S. SOMERVILLE⁴

Institute of Astronomy, Madingley Rd., Cambridge CB3 0HA, UK

Astrophysical Journal Letters, submitted

ABSTRACT

We present a method for constraining the evolution of the galaxy luminosity-velocity (LV) relation in hierarchical scenarios of structure formation. The comoving number density of dark-matter halos with circular velocity of 200 km s^{-1} is predicted in favored CDM cosmologies to be nearly constant over the redshift range $0 \lesssim z \lesssim 5$. Any observed evolution in the density of bright galaxies implies in turn a corresponding evolution in the LV relation. We consider several possible forms of evolution for the zero-point of the LV relation and predict the corresponding evolution in galaxy number density. The Hubble Deep Field suggests a large deficit of bright ($M_V < -19$) galaxies at $1.4 \lesssim z \lesssim 2$. If taken at face value, this implies a dimming of the LV zero-point by roughly 2 magnitudes. Deep, wide-field, near-IR selected surveys will provide more secure measurements to compare with our predictions.

Subject headings: cosmology: theory — dark matter — galaxies: formation — large-scale structure of universe

1. INTRODUCTION

Within the framework of hierarchical structure formation, galaxies form within virialized dark-matter (DM) halos. The halo properties and their time evolution can be predicted robustly within a given Cold Dark Matter (CDM) model using N-body simulations or analytic approximations. A fundamental property of a DM halo is its maximum circular velocity, which is likely to be related to the observable internal velocity of the luminous galaxy that resides in the halo.

In the local universe, it is well known that galaxies obey fundamental scaling relations, including the relation between total luminosity and internal velocity (commonly known as Tully-Fisher or Faber-Jackson relations), which we will focus on here. This LV relationship surely reflects some fundamental aspect of galaxy formation, although its origin is poorly understood. An observational determination of the *redshift evolution* of this relationship would provide crucial clues as to its physical origin and would constitute an important constraint on models of galaxy

formation. However, a direct observational determination of the internal velocities of galaxies at high redshift is extremely difficult. Rotation curves have been obtained for a small number of galaxies up to redshift $z \sim 1$ (Vogt et al. 1996, 1997, 2000). These results suggest that there is little evolution in the Tully-Fisher relation for objects similar to local large spirals out to $z \sim 1$ (but see Mallen-Ornelas et al. 1999). At higher redshifts ($z \gtrsim 1$), no direct observations are yet available.

While galaxies of different morphological types obey different LV relations, one can define a mean LV relation for any given morphological mix. This can be used to make a direct connection between the distribution functions of galaxies in terms of their *internal velocity* and *luminosity*. Clearly, if the evolution of the velocity function is predicted by theory, and the evolution of the luminosity function is determined observationally, this connection can provide a constraint on the evolution of the LV relation.

Determining the number density of galaxies as a function of their rest-frame luminosity is considerably easier than measuring their rotation velocities directly at high

¹james@astronomy.ohio-state.edu

²dekel@astro.huji.ac.il

³joel@ucolick.org

⁴rachel@ast.cam.ac.uk

redshift. Observed luminosities can be k-corrected to the rest visual frame up to $z \sim 2$ using multi-band photometry extending into the near-IR. It has been very difficult to obtain spectroscopic redshifts in the ‘desert’ regime of $1.4 \lesssim z \lesssim 2$, but photometric redshifts offer a means of filling in this gap. These advances, combined with theoretical progress in predicting the number density and structural properties of dark matter halos, suggest the approach for constraining the redshift evolution of the LV relation that we exploit here.

The basis of our analysis is the evolution of the velocity function of DM halos as determined via a high-resolution N-body simulation of the Λ CDM cosmology (Sigad et al. 2000, hereafter S00). We extend these results to other cosmologies using analytic models. We consider a wide range of possibilities for the redshift evolution of the LV relation, including some motivated by the results of semi-analytic models (cf. Somerville & Primack 1999). We then present predictions of the corresponding redshift evolution of the comoving number density of bright galaxies, which may be compared directly with observations to provide quantitative constraints on how the LV relation evolves in time.

2. VELOCITY FUNCTION

We have calculated the velocity function of DM halos using an N-body simulation of the currently popular flat Λ CDM model with $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, Hubble constant $h = 0.7$, and $\sigma_8 = 1.0$. The simulation (Klypin et al. 1999) used a high-resolution adaptive refinement tree code (Kravtsov, Klypin & Khokhlov 1997) to follow the evolution of structure in a periodic box of comoving side $60 h^{-1} \text{Mpc}$, with a force resolution of $\sim 2 h^{-1} \text{kpc}$ in the densest regions.

We identify bound, virialized halos using a spherical overdensity-based algorithm and the standard definition of virial density from the spherical top-hat collapse model (cf. Eke, Cole, & Frenk 1996; Bryan & Norman 1999). Details of the halo finder are described in Bullock et al. (2000; hereafter B00). Although the virial velocity $v_v \equiv [Gm_v/r_v]^{1/2}$ (where m_v and r_v are the virial mass and radius) is often used to approximate galactic circular velocities, a more relevant quantity observationally is the maximum rotation velocity of the halo v_m , which can be related to v_v via the halo density profile. Over relevant radii, halo density profiles are well approximated by the NFW functional form with two free parameters (Navarro, Frenk & White 1996): $\rho(r) = \rho_s(r/r_s)^{-1}(1 + r/r_s)^{-2}$, where r_s is the radius at which the log slope of the profile is -2 . The halo maximum circular velocity occurs at $2.15r_s$ and is related to v_v via $v_m = \sqrt{c/f(c)}v_v$, where $f(c) = 4.62[\ln(1+c) - c/(1+c)]$. Here, c is the ‘concentration’ parameter $c = r_v/r_s$. The value of c is closely related to the halo mass, and for a typical DM halo surrounding a bright galaxy ($m_v \sim 10^{12} h^{-1} M_\odot$) at $z = 0$, our simulations yield $c \sim 13$, implying $v_m = 1.3v_v$. The concentration at a fixed mass evolves rapidly with redshift, $c \propto (1+z)^{-1}$ (B00), implying a corresponding evolution in $v_m(v_v)$. The halo catalogs described in B00 provide v_m and v_v for each halo. A detailed study of the halo velocity

function derived from these catalogs is presented by S00.




FIG. 1.— Evolution of relative comoving number density for fixed $v_m = 200 \text{ km s}^{-1}$ (bold curves) and $v_v = 200 \text{ km s}^{-1}$ halos in three cosmologies.

Figure 1 shows the evolution with redshift of the comoving number density of simulated halos of fixed velocity, either $v_m = 200 \text{ km s}^{-1}$ or $v_v = 200 \text{ km s}^{-1}$. The evolution of halos of fixed v_m is weaker than for halos of fixed v_v due to the strong evolution of c . A fit

$$dn = \psi(v_m) d(\log v_m) = \psi_* v_m^{\tilde{\beta}} d(\log v_m) \quad , \quad (1)$$

for the whole halo v_m distribution is provided by S00, who find that the comoving density at $v_m = 300 \text{ km s}^{-1}$ is constant as a function of z (± 0.1 in the log) at least out to $z = 5$, while $\tilde{\beta} \approx -2.9 - 2.0\tilde{z} + 1.4\tilde{z}^2 \pm 0.1$, where $\tilde{z} \equiv \log(1+z)$.

Figure 1 also shows predictions based on the analytic approximation of Sheth & Tormen (1999, hereafter ST) — a modification of the Press-Schechter approximation (1974) which improves the agreement with simulations. We converted the ST mass to v_v and then to v_m using the analytic model of B00 for $c(m_v, z)$. We see that for Λ CDM, the analytic model provides a good approximation to the simulated evolution both for fixed v_m and fixed v_v . This allows us to extend our predictions using ST to two other cosmological models: τ CDM, a flat model with $\Omega_m = 1$, $h = 0.5$ and a power spectrum of shape parameter $\Gamma = 0.21$, and OCDM, an open geometry with $\Omega_m = 0.3$, $\Omega_\Lambda = 0$, and $h = 0.7$.

In the case of Λ CDM, we note an increase of $\sim 30\%$ in the density for constant v_m between $z = 0$ and $z \sim 3$, and a gradual decline back to today’s density by $z \sim 5$. The density of halos of a constant v_v grows by a factor ~ 2.6 by $z \sim 3$, and then declines at higher z . Note that a halo with a fixed v_v has a decreasing mass and virial radius at increasing redshift. Compared to Λ CDM, the OCDM model predicts a slightly larger increase in the density of

200 km s⁻¹ halos at all redshifts. In the case of τ CDM, the peak density is obtained at $z \sim 1$, followed by a sharper decline, but note that even in this case the comoving density at $z \sim 2$ is comparable to the density at $z = 0$.

3. LV RELATIONS AND LUMINOSITY FUNCTIONS

Armed with this knowledge about the evolution of the halo velocity function, we can now predict the evolution of the number density of galaxies brighter than a fixed absolute luminosity for a given model of LV evolution.

A possible hitch is that the halo maximum velocity v_m typically occurs at a radius r_m that lies outside the luminous parts of the galaxy for which the LV quantities are measured. For example, a $v_m = 200$ km s⁻¹ halo at $z = 0$ has $r_m \sim 45$ kpc. The velocity due to the halo alone obviously declines at smaller radii, but the condensed baryons contribute significantly to the total circular velocity there. We therefore make the assumption that the rotation curve is roughly flat between where it is measured and r_m , i.e. that the halo v_m is the same as the velocity used in the observed LV relations. Gonzalez et al. (1999; G99)⁵ used this assumption at $z = 0$ to compare the velocity function of DM halos with that obtained from observed luminosity functions and LV relations. The good agreement that they obtained is evidence that this is a reasonable approximation at low redshift. We now assume that this relation holds at high redshift as well.

We parameterize the mean LV relation as: $M_B = a(z) - b(\log 2v_m - 2.5) + 5 \log h$, where only the zero-point may evolve in time according to the parametric functional form

$$a(z) = a_0 - 2.5 \log F(z), \quad F(z) = (1 + z^\alpha)^{\beta/\alpha}. \quad (2)$$

We choose $a_0 = -18.71$ and $b = 6.76$, corresponding to the empirical local LV relation used in G99. The parameters α and β allow a wide range of shapes, where α governs the behavior at $z < 1$, and β the asymptotic shape at larger redshifts. The indications for only little evolution at $z \lesssim 1$ guide us to consider mostly large positive values for α , while the value of β is unconstrained, and even its sign is unknown.

In Figure 2 we consider five different representative cases for the LV zero-point evolution $a(z)$, and show the predicted evolution of the number density of galaxies with $M_B \leq -20.2$, relative to their number density at $z = 0$. The case $(\alpha, \beta) = (0, 0)$ is the case of no evolution in the LV relation. The case $(2, -1)$ is close to the evolution predicted for the Λ CDM model if the mass-to-light ratio is constant in time. In this model, because halos of fixed v_m have smaller masses at high redshift, they are fainter and therefore the predicted number of bright galaxies decreases.

The case $(1, 1)$ is close to the prediction of a semi-analytic model (the “accelerated quiescent” model of Somerville, Primack, & Faber 2000) in which the efficiency of star formation scales with the inverse dynamical time of the halo. This scaling is motivated by one of the empirical recipes suggested Kennicutt (1998), based on observations

of nearby galaxies. The dynamical time for a halo of a fixed velocity is smaller at high redshift (because of the higher density), and so galaxies of a fixed velocity are *brighter*. This model predicts an *increase* in the number density of bright galaxies out to $z \sim 5$. Although this star formation recipe is a standard choice in semi-analytic models (e.g., Kauffmann, White, & Guiderdoni 1993), note that it predicts fairly strong evolution in the LV relation even at $z \sim 1$. The other two cases, $(5, 1)$ and $(4, -4)$, are arbitrary choices that illustrate a wide range of possibilities while keeping the LV relation almost unchanged out to $z \sim 1$.




FIG. 2.— Five parameterized models marked by (α, β) for the evolution of the LV zero-point, $a_0 - a(z)$ (Eq. 1; larger values are brighter), along with the implied relative evolution of the comoving density of galaxies brighter than $M_B = -20.2$ for the three cosmological models.

4. DISCUSSION

We have argued that observed estimates of the number density of galaxies above a given luminosity as a function of redshift can be used to constrain the redshift evolution of the LV relation. Lilly et al. (1995) found only little evolution in the number density of *bright* galaxies in the rest-frame B-band out to $z \sim 1$, consistent with the direct constraints on LV evolution out to this redshift (Vogt et al. 1996, 1997, 2000). Redshift surveys of the past decade were unable to reach the higher redshift range $1.0 \lesssim z \lesssim 2.0$ associated with the spectroscopic ‘desert’ or ‘high place of sacrifice’. The results presented by Dickinson (2000), based on NICMOS (near-IR) imaging of the Hubble Deep Field (HDF) suggest that the assembly of present-day luminous galaxies may be taking place precisely in this redshift interval. Using photometric redshifts, Dickinson finds that there are only one-third as many galaxies with rest-frame $M_V < -19$ in the redshift

⁵The published version of this manuscript contains an error in the Equation 12: the quantity in the square root should be inverted.

interval $1.4 \lesssim z \lesssim 2$ as in the interval $0 \lesssim z \lesssim 1.4$. In our fiducial Λ CDM cosmology, the comoving volumes of the HDF for these two intervals are equal, implying an overall drop in the comoving number density at $z \gtrsim 1.4$ of about a factor of three. Taken at face value, this would correspond to dramatic evolution in the LV relation at $z \gtrsim 1$, rather similar to the most extreme ‘dimming’ model shown in Figure 2.

Probably the weakest link in our calculation is our assumption that the halo maximum velocity is the same as the measured internal velocity for a galaxy. The actual measured velocity will depend on the baryon fraction in the galaxy, the spatial distribution of the baryons, and the degree to which angular momentum is preserved in the collapse. However, note that because we expressed the space density in Figure 2 *relative* to that at $z = 0$, our plotted results will not change by much as long as the ratio of observed velocity to the halo maximum velocity, $C \equiv v_{\text{obs}}/v_{\text{m}}$, is of order unity and roughly constant with redshift. We have calculated the value of $C(z)$ using a simple analytic model of disk formation, which assumes that a constant fraction of the halo mass condenses into a disk, and angular momentum is conserved (Mo, Mao, & White 1998). We find that $C(z) \simeq 1.3$ for the redshift range $0 \lesssim z \lesssim 2$, and falls to $C(z) \simeq 1.0$ by $z \sim 5$. This suggests that our predictions should be reasonably robust at least out to $z \sim 2$.

How significant is the effect seen in the HDF? One possibility is that galaxies are preferentially missed at $z \gtrsim 1.4$ because of the $(1+z)^4$ cosmological surface-brightness dimming. Dickinson (2000) claims that this is unlikely to be a large effect; when bright HDF galaxies from $z \lesssim 1$ are artificially redshifted to $z \lesssim 2$ with no intrinsic luminosity evolution, most of them are still easily detectable. Another possibility is that the photometric redshifts are inaccurate; Dickinson points out that there are *no* spectroscopic redshifts in the range $1.4 \lesssim z \lesssim 2$ where the deficit is seen. However, this would require quite a dramatic fail-

ure where *all* of the galaxies (with photometric redshifts) assigned to $0.5 \lesssim z_{\text{phot}} \lesssim 1.4$ would have to be reassigned to the higher redshift range. Most importantly, recall that the HDF covers a relatively small volume, and therefore random fluctuations along the line of sight are likely to be large. The observed deficit of galaxies might simply reflect large-scale ‘spikes’ in the redshift distribution. Although there is some evidence for a similar deficit of bright galaxies at $z \gtrsim 1$ in other deep optical/IR surveys (Fontana et al. 1999), secure confirmation of these results will require deep wide-field surveys with multi-band optical-IR photometry. Several such surveys are now in progress (e.g. the Cambridge-Carnegie Las Campanas IR Survey). Moreover, the DEEP survey (Davis & Faber 1998) will obtain measured line-widths for a large number of galaxies at $z \gtrsim 0.7$, which will provide direct constraints on the LV relation at high redshift.

Finally, our results may be relevant to a recent proposal by Newman & Davis (2000) to use velocity-selected galaxies as a tracer population in a classical dN/dz measure of cosmological parameters. This analysis would be subject to similar uncertainties in relating the halo velocity to observable galaxy velocity, e.g., the dependence on whether the halos are selected by v_{m} or v_{v} . Moreover, because the proposed spectroscopic surveys are magnitude limited, the indicated evolution of the LV relation could introduce severe selection biases into the analysis.

ACKNOWLEDGMENTS

This work has been performed at the ITP, Santa Barbara. It has been supported by grants from the US-Israel BSF, the Israel SF, NASA and NSF at UCSC and NMSU. JSB was supported by NASA LTSA grant NAG5-3525 and NSF grant AST-9802568. JRP acknowledges a Humboldt Award at the Max Planck Institute for Physics in Munich. RSS is supported by a rolling grant from PPARC. We thank Marc Davis and Mark Dickinson for helpful correspondence.

REFERENCES

- Bryan, G.L., Norman, M.L. 1998, ApJ, 495, 80
 Bullock, J.S., Kolatt, T.S., Sigad, Y., Somerville, R.S., Kravtsov, A.V., Klypin, A.A., Primack, J.R., & Dekel, A. 2000, MNRAS submitted, astro-ph/9908159 (B00)
 Davis, M. & Faber, S.M. 1998, in ‘Wide Field Surveys in Cosmology’, eds. S. Colombi & Y. Mellier, astro-ph/9810489
 Dickinson, M., 2000, in ‘Building Galaxies: From the Primordial Universe to the Present’, eds. Hammer, F., Thuan, T.X., Cayatte, V., Guiderdoni, B., & Tranh Than Van J. (Paris:Ed. Frontières), p. 257, astro-ph/0004027
 Eke, V.R., Cole, S., Frenk, C.S. 1996, MNRAS, 282, 263
 Fontana, A., et al. 1999, MNRAS, 310, L27
 Gonzalez, A.H., Williams, K.A., Bullock, J.S., Kolatt, T.S., & Primack, J.R. 2000, ApJ, 528, 145
 Kauffmann, G., White, S.D.M., & Guiderdoni, B., 1993, MNRAS, 264, 201
 Kennicutt, R. 1998, ApJ, 498, 541
 Klypin, A.A., Gottlöber, S., Kravtsov, A.V., & Khokhlov, A.M. 1999, ApJ, 516, 530
 Kravtsov, A., Klypin, A., & Khokhlov, A.M. 1997, ApJS, 111, 73
 Lilly, S.J., Tresse, L., Hammer, F., Crampton, D., & Le Fèvre, O., 1995, ApJ, 455, 108
 Mallen-Ornelas, G., Lilly, S.J., Crampton, D., & Schade, D. 1999, ApJL, 518, 83
 Navarro, J., Frenk, C., & White S.D.M. 1996, ApJ, 462, 563
 Newman, J.A. & Davis, M., 2000, ApJL, accepted, astro-ph/9912366
 Press, W. & Schechter, P. 1974, ApJ, 187, 425
 Sheth, R.K., & Tormen, G. 1999, MNRAS, 308, 119 (ST)
 Sigad, Y., Kolatt, T.S., Bullock, J.S., Kravtsov, A.V., Klypin, A.A., Primack, J.R., & Dekel, 2000, MNRAS, submitted (S00)
 Somerville, R.S. & Primack, J.R. 1999, MNRAS, 310, 1087
 Somerville, R.S., Primack, J.R., & Faber, S.M., 2000, MNRAS, submitted
 Vogt, N.P., et al., 1996, ApJ, 465, 15
 Vogt, N.P., et al., 1997, ApJ, 479, L121
 Vogt, N.P., et al., 2000, in prep.